



A trans-Atlantic examination of haddock *Melanogrammus aeglefinus* food habits

Tam, J.C.; Link, J.S.; Large, S.I.; Bogstad, B.; Bundy, A.; Cook, A.M.; Dingsør, G.E.; Dolgov, A.V.; Howell, D.; Kempf, A.

Total number of authors:
15

Published in:
Journal of Fish Biology

Link to article, DOI:
[0.1111/jfb.12983](https://doi.org/10.1111/jfb.12983)

Publication date:
2016

Document Version
Peer reviewed version

[Link back to DTU Orbit](#)

Citation (APA):

Tam, J. C., Link, J. S., Large, S. I., Bogstad, B., Bundy, A., Cook, A. M., Dingsør, G. E., Dolgov, A. V., Howell, D., Kempf, A., Pinnegar, J. K., Rindorf, A., Schückel, S., Sell, A. F., & Smith, B. E. (2016). A trans-Atlantic examination of haddock *Melanogrammus aeglefinus* food habits. *Journal of Fish Biology*, 88(6), 2203-2218. <https://doi.org/10.1111/jfb.12983>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

1 TITLE: A Trans-Atlantic Examination of Haddock (*Melanogrammus aeglefinus*) Food Habits

2
3 AUTHORS: J. C. Tam^{1*}, J. S. Link¹, S. I. Large^{1,2}, B. Bogstad³, A. Bundy⁴, A. M. Cook⁴, G. E.
4 Dingsør³, A. Dolgov⁵, D. Howell³, A. Kempf⁶, J. K. Pinnegar⁷, A. Rindorf⁸, S. Schüeckel⁹, A. F.
5 Sell⁶, B. E. Smith¹

6
7 ADDRESSES:

8 ¹NOAA-Fisheries, 166 Water Street, Woods Hole, MA 02543, USA

9 ²International Council for the Exploration of the Seas (ICES), Copenhagen V 1553, Denmark

10 ³Institute of Marine Research (IMR), 5817, Bergen, Norway

11 ⁴Fisheries and Oceans Canada, Bedford Institute of Oceanography, Dartmouth, NS, B2Y 4A2,
12 Canada

13 ⁵Polar Research Institute of Marine Fisheries and Oceanography (PINRO), 6, Knipovich-St.,
14 Murmansk, 183038, Russia

15 ⁶Thünen Institute of Sea Fisheries, Palmaille 9, 22767 Hamburg, Germany

16 ⁷Centre for Environment, Fisheries & Aquaculture Science (Cefas), Pakefield Road, Lowestoft,
17 Suffolk, NR33 0HT, UK

18 ⁸National Institute of Aquatic Resources, Technical University of Denmark, Charlottenlund Slot,
19 DK-2920 Charlottenlund, Denmark

20 ⁹BioConsult, Reeder-Bischoff Str. 54, 28757 Bremen, Germany

21 *Corresponding author: jamie.tam@noaa.gov +01 508 495 2083

22
23 RUNNING HEADLINE: Trans-Atlantic examination of haddock diet

ABSTRACT

Haddock (*Melanogrammus aeglefinus*) is a commercially and ecologically important gadoid in the food webs and economies of regional ecosystems across the North Atlantic. *Melanogrammus aeglefinus* is reported to be primarily piscivorous in some cases and primarily an echinoderm feeder in others. Often, the species exhibits ontogenetic dietary shifts, focusing on a specific prey at small sizes and on different prey at larger sizes. Here we explore and contrast the food habits of *M. aeglefinus* across multiple northeastern and northwestern Atlantic ecosystems, using databases that span multiple decades. The results show that, among all ecosystems, echinoderms are a consistent part of *M. aeglefinus* diet, but patterns do emerge regarding where and when *M. aeglefinus* primarily eat fishes versus echinoderms. *Melanogrammus aeglefinus* does not regularly exhibit the increase in piscivory with ontogeny that other gadoids often show, in several ecosystems there is a lower occurrence of piscivory. There is an apparent inverse relationship between the consumption of fishes and echinoderms in *M. aeglefinus* over time, where certain years show high levels of one prey item and low levels of another. We view this apparent binary choice as part of a gradient of prey options, contingent upon a suite of factors external to *M. aeglefinus* dynamics. We discuss the energetic consequences of this prey choice, noting that in some instances it may not be a choice at all.

KEY WORDS: haddock, prey choice, echinoderms, piscivory, ecosystem-based fisheries management

INTRODUCTION

An integral part of managing commercial fish stocks from an Ecosystems-Based Fisheries Management (EBFM) perspective is understanding trophic linkages and the nature of feeding choices in target species (Pauly *et al.*, 1998; Link *et al.*, 2002). For predatory species, prey selection defines their ecological role and offers insight into how these species can and should be managed (Greenstreet *et al.*, 1997; Link *et al.*, 2009; Branch *et al.*, 2010). For historically harvested fish stocks, such as gadoids and pleuronectiformes in the North Atlantic, it is important to understand their functional roles and what ecological buffers exist (in terms of functional redundancies, where multiple species contribute in equal ways to ecosystem function) against direct and indirect effects of overexploitation (Bellwood *et al.*, 2003; Link, 2007). In the case of geographically-widespread, opportunistic predators, populations of the same species can develop differing dietary strategies and specializations. This has been well documented in marine mammals where species such as harbour seals *Phoca vitulina* and killer whales *Orcinus orca* exhibit a variety of prey specializations between their geographically widespread populations even though they are able to eat a substantial range of prey (Ford *et al.*, 1998, 2010; Tollit *et al.*, 1998). Fewer such comparisons are made between fish populations, but examinations of fisheries literature indicates that dietary specializations occur in opportunistic fish predators. For example, the spiny dogfish *Squalus acanthias* is a widespread commercial fish that appears to consume proportionately more benthic invertebrates in New Zealand (Hanchet, 1991), teleost fishes in the North Atlantic (Holden, 1966; Link *et al.*, 2002) and large proportions of Euphausiids in the Eastern Pacific (Jones & Geen, 1977; Tanasichuk *et al.*, 1991). This can have strong implications on how target species are managed in different parts of the world as local

abundances of preferred prey can dictate the stock size of the predator and subsequent functional redundancies may differ from place to place (ecosystem to ecosystem).

Haddock *Melanogrammus aeglefinus* is an important species in the North Atlantic, with a long history of commercial exploitation (Jensen, 1965; Bergstad *et al.*, 1987; Pope & Macer, 1996; Hedger *et al.*, 2004). *Melanogrammus aeglefinus* is a widespread species with fisheries in the east coast of North America and in many parts of northern Europe and the British Isles (Aleev, 1944; Jensen, 1965; Albert, 1994; Jiang & Jørgensen, 1996; Temming *et al.*, 2004; Bogstad *et al.*, 2013). Juvenile *M. aeglefinus* transition after 3-5 months from living in the upper reaches of the water column to living on muddy, sandy or gravel substrate (Platt *et al.*, 2003; Sell & Kröncke, 2013). As demersal fish, they spend their adult life in close association with benthic habitat. Like other gadoids, they are opportunistic, higher-trophic-level feeders and prey on a variety of fishes and invertebrates (Schückel *et al.*, 2010; Smith & Link, 2010). Although *M. aeglefinus* is an abundant species in many ecosystems and its feeding behaviour is extensively studied in specific locales (Tseeb, 1964; Sonina, 1969; Dolgov *et al.*, 2007), overviews about its diet or how its diet may change throughout its life history or across its geographic range have not been executed comparatively.

Echinoderms have been relatively well studied in marine ecology, primarily as consumers. Notable examples are crown-of thorns *Acanthaster planci* decimating corals in the Indo-Pacific (Moran *et al.*, 1992; Vogler *et al.*, 2008), seastars *Pisaster ochraceus* as a keystone predator on North Pacific intertidal communities (Paine, 1966, 1969) and sea urchins *Strongylocentrotus droebachiensis* denuding temperate areas entirely of macrophytes (Watanabe & Harrold, 1991;

Gagnon *et al.*, 2004; Lauzon-Guay & Scheibling, 2007). However, outside of the sea otter-urchin-macrophyte feeding dynamic (Estes & Palmisano, 1974; Dayton, 1975), studies of predation on echinoderms seldom occur (but see Pinnegar *et al.*, 2000; Schückel *et al.*, 2010). Furthermore, the trophic role of echinoderms in deeper, subtidal communities is undervalued commercially as well as ecologically and not well studied compared to the role of fish prey, particularly in the northern Atlantic.

A unique aspect of *M. aeglefinus* feeding is that echinoderms can make up a significant part of their diet in the Northwestern Atlantic (Mahon & Neilson, 1987; Link, 2004; Shackell *et al.*, 2010; Smith & Link, 2010). This is rare among gadoids and from an energetic perspective would appear to be counterintuitive (Dauvin & Joncourt, 1989). With the exception of sea cucumbers, echinoderms are mainly composed of thick, calcareous exoskeletons that are only slowly digested by most fishes and contain less energy per gram than other invertebrate diets items including worms, gastropods, bivalves and crustaceans (Steimle & Terranova, 1980). Survey observations of fish stomach contents in the Northwest Atlantic has shown that several species consume echinoderms at some point in their life histories (Link, 2004; Smith & Link, 2010; Bundy *et al.*, 2011). The fact that commercially important fishes, like *M. aeglefinus*, choose to eat echinoderms is rarely addressed in fisheries management and ecosystem modelling even though echinoderms appear to play a large role in regulating energy flow in marine food webs (Steimle & Terranova, 1980; Zamarro, 1992).

What is unclear is the level of consistency across *M. aeglefinus* populations throughout their range in consuming echinoderms as a significant food source. There are indications that the diet

of *M. aeglefinus* differs between areas and time periods. As a consequence, the functional redundancies to *M. aeglefinus* remain unknown. This study aims to examine the diet similarities of *M. aeglefinus* populations throughout their distributional range, at different stages during their life history and across several decades. This is an important step in understanding the dynamics in feeding ecology of *M. aeglefinus* and the role of echinoderms in benthic ecosystems.

MATERIALS AND METHODS

Study areas and data collection

M. aeglefinus stomachs were collected from nine ecosystems in the North Atlantic (Figure 1). In the Gulf of Maine, Georges Bank, Scotian Shelf, Rockall Bank, North Sea and Barents Sea diet composition was assessed as the weight of the diet item relative to the total weight of the stomach contents. In the Celtic Sea, Irish Sea and Spitzbergen (Svalbard), frequency of occurrence (FO; the number of stomachs containing the diet item relative to the total number of stomachs) was used to assess the diet (Hyslop E. J., 1980; Ahlbeck *et al.*, 2012). The sampling primarily occurred from the 1970s to present, but extended to earlier decades for some regions (Table 1).

Stomach contents were collected from length-stratified samples during fishery-independent and fishery-dependent surveys, and prey were quantified by weight or number depending on where and what type sampling occurred (Table I). Where possible, prey items were

identified to species level; however, due to varying levels of digestion prey were often identified to genus, class or family levels (Smith & Link, 2010; Bundy *et al.*, 2011; Cook & Bundy, 2012).

Data analysis

Data were parsed to include only the dominant prey of *M. aeglefinus* diet. Because they have been described elsewhere in detail for given ecosystems (Table I), non-echinoderm invertebrates (worms, gastropods, bivalves and crustaceans) were not considered in the analyses. As each ecosystem may contain a different selection of prey species and given the aims of this study, we grouped prey according to rough taxonomic classification of echinoderm, fish, or other prey item. To determine if *M. aeglefinus* undergo similar ontogenetic shifts in diet composition across regions, we also grouped *M. aeglefinus* into the following size groups: 10-29, 30-49, and > 50 cm where data were available. These sizes are consistent with observed major ontogenetic shifts in diet (Link & Garrison, 2002a). Where time series data was available (Georges Bank, Gulf of Maine, US Scotian Shelf and Barents Sea), *M. aeglefinus* diet was plotted in 5 year blocks to examine consistency in food choice over time.

Given that the diet of *M. aeglefinus* was recorded differently between ecosystems, Kendall's tau (Fredricks & Nelsen, 2007) was used to examine the non-parametric, rank-order relationship between the proportion of echinoderm and fish in the diet of *M. aeglefinus* in differing ecosystems. This statistical method is the non-parametric equivalent of measures of association or correlation. This analysis determined if there was any consistent and significant association between the amount of fishes and echinoderms in the diet of *M. aeglefinus* for all

ecosystems examined. Within ecosystem differences between echinoderms and fishes in *M. aeglefinus* diet were determined using Chi-square analysis. To explore and test for relationships within ecosystems of *M. aeglefinus* diet between size classes, Chi-square analysis of the proportion of each diet item in the stomachs of fish across size classes was used. To examine trends in prey selection of *M. aeglefinus* at each ecosystem over time, a generalized linear model was used with percent of the diet item as the dependent variable and time as the independent variable. This determined if *M. aeglefinus* showed general shifts in their prey selection over time. At each 5 year block within each ecosystem, a Chi-square analysis was performed to determine significant differences between the percentage of echinoderms and fishes in *M. aeglefinus* diet and reveal any changes in the feeding pattern of *M. aeglefinus* over time.

RESULTS

There was no significant relationship of echinoderms and fishes in the diet of *M. aeglefinus* that could be applied to all ecosystems according to the Kendall's tau analysis (Figure 2; $\tau = 0.33$, Z-score = 1.25, $p = 0.21$). The North Sea and Rockall Bank populations of *M. aeglefinus* showed significantly higher piscivory, while all other ecosystems showed significantly higher echinoderms consumed or else no dominant prey (Table II).

Differing patterns of ontogenetic shifts towards piscivory were apparent in *M. aeglefinus* among ecosystems based on size class data (Figure 3). In several ecosystems there was a significantly lower occurrence of piscivory (i.e. eggs and larvae) with increasing size (Georges Bank: $X^2 = 30.52$, $df = 2$, $p = <0.001$; Gulf of Maine: $X^2 = 15.91$, $df = 2$, $p = <0.001$). In the

North Sea and Rockall Bank there was a higher occurrence of piscivory (i.e. small pelagic fishes) with increasing size class ($X^2 = 38.34, 23.90, df = 2, 2, p < 0.001, <0.001$, respectively), but the proportion of echinoderms in the diet of *M. aeglefinus* remained constant. Through their life history, *M. aeglefinus* ate a large proportion of echinoderms in most ecosystems. Echinoderms were a proportionately consistent component of *M. aeglefinus* diet throughout their life cycle except in the Gulf of Maine and Scotian Shelf, where echinoderms increased in frequency with increasing size ($X^2 = 22.29, 17.82, df = 2, 2, p < 0.001, <0.001$, respectively).

There was no consistent temporal shift in the diet of *M. aeglefinus* in either echinoderms or fishes within ecosystems ($F_{1,4} = 0.18, p = 0.95$). Long-term stomach content data indicated that echinoderms were consistently a part of *M. aeglefinus* diet over time, but neither diet item exhibited any positive or negative linear trend, indicating that there was no shift in diet related to an external pressure (i.e. environment or habitat). Despite the lack of trend, the Chi-square analyses indicated that there were significant differences in the consumption of echinoderms and fishes by *M. aeglefinus* within time periods (Table III). There appeared to be an inverse relationship between the consumption of fishes and echinoderms by *M. aeglefinus* over time. Periods of high and low fish consumption by *M. aeglefinus* coincided with respective periods of low and high echinoderm consumption (Figure 4). During the late 1980s and early 2000s there appeared to be peaks in echinoderm consumption by *M. aeglefinus*.

DISCUSSION

It is unusual for a gadoid species to consistently maintain a high proportion of benthic invertebrates in its diet throughout its life history. Most gadoids are known to be opportunistic:

Atlantic cod *Gadus morhua* being a generalist feeder, saithe *Pollachius virens* and some hake species consuming more of a pelagic diet (Smith & Link, 2010). *Gadus morhua* have shown a clear ontogenetic shift in diet, with smaller *G. morhua* feeding more on benthic prey and larger *G. morhua* showing a strong shift towards piscivory (Link & Garrison, 2002b; Smith *et al.*, 2007; Rowlands *et al.*, 2008). In contrast, *M. aeglefinus* exhibit no ontogenetic shift towards piscivory with the exception of the North Sea and Rockall Bank, where echinoderms still occur in their diets throughout their lives. This difference in ontogenetic diet shift between *M. aeglefinus* and *G. morhua* can be explained by the smaller gape size and body size of *M. aeglefinus* that are not able to eat as many larger prey fish compared to *G. morhua* (Løkkeborg *et al.*, 1989).

M. aeglefinus consistently consume echinoderms across the range of environmental conditions examined. In ecosystems where *M. aeglefinus* appear to be more piscivorous or predominantly eat a combination of other invertebrates, echinoderms still make up at least 10% of their diet. Since the early 2000s *M. aeglefinus* stocks have experienced an increase in abundance in all regions studied with the exception of the Scotian Shelf and, thus, do not appear to be suffering from reduced productivity due to their reliance on low-nutrient food sources (Brodziak *et al.*, 2008; Bogstad *et al.*, 2013; Pecuchet *et al.*, 2014). Eating food of high calcium carbonate and low energy may have developed in *M. aeglefinus* to reduce competition with other gadoids by obtaining a consistent, albeit low-energy food source.

While it may not seem like a successful strategy to consume large proportions of echinoderms due to their low nutritional value compared to prey fish, their relatively high

abundances and relatively low defenses (slow moving) make them a very accessible prey to *M. aeglefinus*. Echinoderms in the northwestern Atlantic make up a large proportion of the biomass in benthic habitats (up to 500 g m⁻² and up to 90% of the total grab sample) with sand dollars *Echinarachinius parma* in shallower regions and brittlestars *Amphioplus* spp. and sea cucumbers dominating deeper areas of Georges Bank and the Gulf of Maine basins (Steimle, 1990). In Atlantic Canada, echinoderms at times have made up to 35% of the total cal m⁻² of benthic habitat even though they are relatively low in energy content compared to other invertebrate phyla (Brawn *et al.*, 1968). In the North Sea, the brittlestar *Amphiura filiformis* dominates the benthic system and lives in densities up to 2250 indv m⁻² (Skold *et al.*, 1994; Wood *et al.*, 2009), and other important species include *Echinocyamus pusillus*, *Ophiura albida*, *Ophiura affinis*) and *Echinocardium cordatum* that are also common components of the benthos. These high abundances of echinoderms in benthic communities support the hypothesis that *M. aeglefinus* exhibit ambient feeding in some regions to perhaps sustain the populations while supplementing their diet with other prey.

In this study the diet of *M. aeglefinus*, consistently include echinoderms across ecosystems and over time, even in regions where the dominant prey is fish. This indicates that echinoderms are an important food source throughout their distribution. While there does not appear to be long-term, multi-decadal shifts in the diet of *M. aeglefinus*, there is an indication of shorter-term changes in *M. aeglefinus* feeding habits (Figure 4). At five year intervals in Georges Bank, Gulf of Maine, Scotian Shelf and Barents Sea there is evidence of an inverse relationship between the contribution of echinoderms and fish prey to the diet of *M. aeglefinus* (Antipova *et al.*, 1980). Foraging fish are known to be able to alter their behaviour in response to

variations in food availability and the threat of competition (Dill, 1983). *Melanogrammus aeglefinus* likely consume lower quality food choices when faced with adversity in acquiring a preferred resource (Vinyard, 1980; McNamara & Houston, 1987). Feeding on echinoderms could be a strategy for *M. aeglefinus* to maintain constant energy levels during periods of low prey fish abundance and high competition with other gadoids that may occur on a multi-annual cycle (Temming *et al.*, 2004). However, intra-annual abundances for all prey items of *M. aeglefinus* from the ecosystems surveyed in this study were not examined and it would be a useful avenue for future research to measure selectivity across time in all of these populations.

An opportunistic feeding mode for *M. aeglefinus* has been demonstrated in several earlier studies (Mattson, 1992; Schückel *et al.*, 2010) and current observations from the different ecosystems reveals that high amounts of fish in the diet of larger North Sea *M. aeglefinus* is likely to be a consequence of prey availability rather than an ontogenetic shift (i.e. the high abundance of sandeels *Ammodytes* spp.). This is supported by observations on Dogger Bank in the North Sea, where Sell and Kröncke (2013) found the distribution of *M. aeglefinus* to be more strongly correlated with abiotic habitat parameters (depth, temperature, sediment composition) than with biotic factors, including the availability of potential mud dwelling prey. It has also been shown that *M. aeglefinus* adapt their diets based on prey availability due to benthic disturbance from fishing gear with less benthos and slightly more fish consumed in disturbed habitats (Smith *et al.*, 2013).

There is some indication that echinoderms have higher nutrition and energy content than expected, but past studies have shown that the caloric content of fish to be higher than

echinoderms. Brawn et al. (1968) found that prey fish contained 2-3 times more cal g⁻¹ than echinoderms in St. Margaret's Bay in Nova Scotia, Canada. Other studies have found lower energy content in seastars and brittlestars compared to fish (19.10 kJ g⁻¹ for echinoderms and 22.96-24.22 kJ g⁻¹ for fish; ash-free dried weight), while sea urchins had a higher energy content (25.82 kJ g⁻¹; ash-free dried weight) than both demersal or pelagic fishes (Steimle & Terranova, 1980; Packer *et al.*, 1994). Inconsistencies in the values of energy contents drawn from these studies may be due to the different species contributing to the sample groups. There may also be a seasonal element to the appeal of echinoderms as *M. aeglefinus* prey. When echinoderms are gravid they are more nutritionally valuable and contain a higher lipid and protein content. Steimle and Terranova (1980) found that the energy content in *E. parma* doubled during autumn spawning periods compared to spring values. Certain conditions in timing and availability may make echinoderms a logical food choice for *M. aeglefinus*.

There is some evidence that hard shelled invertebrates, such as echinoderms and bivalves, digest more slowly and less efficiently than other, soft-tissue prey such as fishes and polychaetes (MacDonald *et al.*, 1982; Tibbetts *et al.*, 2004; Jaworski & Ragnarsson, 2006). Preliminary calculations indicate that the digestibility of hard shelled invertebrates are only approximately 10% less than that of soft bodied prey such as fish and polychaetes in gadoid diets, and respective evacuation rates could be on the order of 1-2 times slower (MacDonald *et al.*, 1982; Tibbetts *et al.*, 2006). If this is the case, the relative importance of echinoderms in the diet of *M. aeglefinus* could potentially be amplified by a similar factor. If, however, the percent diet composition of echinoderms in the diet of *M. aeglefinus* is lower, the fact remains that echinoderms still remain a notable, consistent and regular component of the diet of *M.*

aeglefinus. In contrast, there is evidence that the shells in echinoderms may actually promote more efficient digestion in predatory fish. Mattson (1992) indicates that the calcareous shell of echinoderms act as a grinding element and can enhance digestion as well as ingestion. This increased maceration may compensate for the low nutritional value inherent in echinoderm prey and could ultimately improve the digestion of all hard shelled organism in *M. aeglefinus* diet.

The implications of understanding the diets of target fish species span far beyond just knowing what they are eating. Assessing food habits in fishes, particularly in opportunistic feeders, can give insight into habitats and prey species distributions that are difficult to survey (Link, 2004; Cook & Bundy, 2012). In determining that *M. aeglefinus* consume large proportions of echinoderms in many regions, this study has identified an important link between the benthos and harvested fish. This is a mechanism that is rarely considered in management decisions and implies that impacts to the benthic habitat in certain regions will likely affect *M. aeglefinus* more than other harvested gadoid species. From an EBFM perspective, the functional redundancies of *M. aeglefinus* lie alongside other echinoderm consumers such as American plaice *Hippoglossoides platessoides* and ocean pout *Macrozoarces americanus* rather than species of the same family group (Zamarro, 1992; Packer *et al.*, 1994; Link, 2007).

While traditional conservation strategies, such as fisheries closures, are not always effective at increasing target fish stocks (Fisher & Frank, 2002; Kelly *et al.*, 2006), *M. aeglefinus* have historically responded well to these management options (Brodziak *et al.*, 2008). In the mid-1990s, *M. aeglefinus* stocks in the northwest Atlantic were the lowest on record (Hutchings, 2000). After strict closures as a response by the declining stock status, *M. aeglefinus* stocks

increased dramatically (Murawski *et al.*, 2000; Brodziak *et al.*, 2008). *Melanogrammus aeglefinus* are also known to have a restricted daily ambit compared to other gadoids, indicating a strong association with local habitats (Perry & Smith, 1994; Fogarty *et al.*, 2001). Reduced impacts to the benthic habitat compounded with the high proportion of echinoderms in *M. aeglefinus* diet, in part, explain why *M. aeglefinus* stocks have successfully responded to fisheries closures and MPAs (Smith *et al.*, 2013). In areas where *M. aeglefinus* exhibited a high frequency of piscivory, such as the North Sea, fisheries closures did not improve their abundances nearly as much (FSBI, 2001; Fisher & Frank, 2002). These differences in feeding habits between populations can complicate management efforts because species may not respond the same way in all regions of their distributional range. Thus, conservation methods need to take into account the biology of the ecosystem as well as species-specific life histories to ensure positive and efficient management outcomes.

The importance of the benthic habitat to harvested fish is becoming increasingly apparent. The interconnectedness and complexity of food webs is an important consideration in sustainable fisheries management (Pauly *et al.*, 1998; Link, 2011; Cook & Bundy, 2012; Link & Auster, 2013). To complicate our understanding of trophic webs in fisheries even further, comparative studies such as this one are revealing that populations of the same fish species can interact with their local habitat differently between regions. This emphasizes the need for ecosystem-level management that will improve the resiliency and long-term economic viability of harvested fish throughout their distributions.

ACKNOWLEDGEMENTS

We would like to thank the research vessel crews at NOAA and DFO whose hard work has made studies such as this possible. We also thank ICES and IMARES for making this stomach data available. Partial support for data collection was provided by EU FP7 project BENTHIS (KBBE.2012.1.2-09) (J.K.P.). We would like to thank internal reviewers Kristin Holsman, Ron Heintz and Kenric Osgood for their helpful comments and suggestions. This work was supported by a NOAA Postdoctoral Fellowship to JCT.

REFERENCES

- Ahlbeck, I., Hansson, S. & Hjerne, O. (2012) Evaluating fish diet analysis methods by individual-based modelling. *Canadian Journal of Fisheries and Aquatic Sciences*, **69**, 1184–1201.
- Albert, O.T. (1994) Ecology of haddock (*Melanogrammus aeglefinus* L.) in the Norwegian Deep. *ICES Journal of Marine Science*, **51**, 31–44.
- Aleev, V.R. (1944) Haddock of the Barents Sea. Fishery and biology review. *PINRO Proceedings*, 187–233.
- Antipova, T.V.I., Ponomarenko, Y.A. & Yaragina, N.A. (1980) Seasonal and year-to-year dynamics of occurrence frequency of the main food objects for haddock and their stomach fullness degree in the Barents Sea. *ICES Council Meeting*.
- Bellwood, D.R., Hoey, A.S. & Choat, J.H. (2003) Limited functional redundancy in high diversity systems: resilience and ecosystem function on coral reefs. *Ecology Letters*, **6**, 281–285.
- Bergstad, O.A., Jorgensen, T. & Dragesund, O. (1987) Life History and Ecology of the Gadoid Resources of the Barents Sea. *Fisheries Research*, **5**, 119–161.

366 Bogstad, B., Dingsør, G.E., Ingvaldsen, R.B. & Gjøsæter, H. (2013) Changes in the relationship
 367 between sea temperature and recruitment of cod, haddock and herring in the Barents Sea. *Marine*
 368 *Biology Research*, **9**, 895–907.

369 Branch, T.A., Watson, R., Fulton, E.A., Jennings, S., McGilliard, C.R., Pablico, G.T., *et al.*
 370 (2010) The trophic fingerprint of marine fisheries. *Nature*, **468**, 431–5.

371 Brawn, V.M., Peer, D.L. & Bentley, R.J. (1968) Caloric content of the standing crop of benthic
 372 and epibenthic invertebrates of St. Margaret's Bay, Nova Scotia. *Journal of Fisheries Research*,
 373 **25**, 1803–1811.

374 Brodziak, J., Traver, M.L. & Col, L.A. (2008) The nascent recovery of the Georges Bank
 375 haddock stock. *Fisheries Research*, **94**, 123–132.

376 Bundy, A., Link, J.S., Smith, B.E. & Cook, A.M. (2011) You are what you eat, whenever or
 377 wherever you eat it: an integrative analysis of fish food habits in Canadian and U.S.A. waters.
 378 *Journal of Fish Biology*, **78**, 514–39.

379 Cook, A.M. & Bundy, A. (2010) *The food habits database: an update, determination of*
 380 *sampling adequacy and estimation of diet for key species.*

381 Cook, A.M. & Bundy, A. (2012) Spies of the ocean: improving our understanding of biodiversity
 382 and ecosystem functioning using fish as sampling tools. *Marine Ecology Progress Series*, **454**,
 383 1–18.

384 Dauvin, J.-C. & Joncourt, M. (1989) Energy values of marine benthic invertebrates from the
 385 Western English Channel. *Journal of the Marine Biological Association of the UK*, **69**, 589–595.

386 Dayton, P.K. (1975) Experimental studies of algal canopy interactions in a sea otter-dominated
 387 kelp community at Amchitka Island, Alaska. *Fisheries Bulletin*, **73**, 230–237.

388 Dill, L. (1983) Adaptive flexibility in the foraging behaviour of fishes'. *Canadian Journal of*
389 *Fish and Aquatic Sciences*, **40**, 398–408.

390 Dolgov, A. V., Yaragina, N.A., Orlova, E.L., Bogstad, B., Johannesen, E. & Mehl, S. (2007)
391 20th anniversary of the PINRO-IMR cooperation in the investigations of fish feeding in the
392 Barents Sea – results and perspectives. In *Proceedings of the 12th Norwegian-Russian*
393 *Symposium* (ed. by Haug, T., Misund, O.A., Gjosaeter, H. & Rottingen, I.). Tromso, pp. 44–78.

394 Estes, J.A. & Palmisano, J.F. (1974) Sea Otters: Thir Role in Structuring Nearshore
395 Communities. *Science*, **185**, 1058–1060.

396 Fisher, J.A.D. & Frank, K.T. (2002) Changes in finfish community structure associated with an
397 offshore fishery closed area on the Scotian Shelf. *Marine Ecology Progress Series*, **240**, 249–
398 265.

399 Fisheries Society of the British Isles (FSBI). (2001) *Marine Protected Areas in the North Sea*
400 *Briefing Paper 1*. Cambridge, UK.

401 Fogarty, M.J., Myers, R.A. & Bowen, K.G. (2001) Recruitment of cod and haddock in the North
402 Atlantic: a comparative analysis. *ICES Journal of Marine Science*, **58**, 952–961.

403 Ford, J.K.B., Ellis, G.M., Barrett-Lennard, L.G., Morton, A.B., Palm, R.S. & Balcomb III, K.C.
404 (1998) Dietary specialization in two sympatric populations of killer whales (*Orcinus orca*) in
405 coastal British Columbia and adjacent waters. *Canadian Journal of Zoology*, **76**, 1456–1471.

406 Ford, J.K.B., Ellis, G.M., Olesiuk, P.F. & Balcomb, K.C. (2010) Linking killer whale survival
407 and prey abundance: food limitation in the oceans' apex predator? *Biology Letters*, **6**, 139–142.

408 Fredricks, G. a. & Nelsen, R.B. (2007) On the relationship between Spearman's rho and
409 Kendall's tau for pairs of continuous random variables. *Journal of Statistical Planning and*

410 *Inference*, **137**, 2143–2150.

411 Gagnon, P., Himmelman, J.H. & Johnson, L.E. (2004) Temporal variation in community
 412 interfaces: kelp-bed boundary dynamics adjacent to persistent urchin barrens. *Marine Biology*,
 413 **144**, 1191–1203.

414 Greenstreet, S.P.R., Bryant, A.D., Broekhuizen, N., Hall, S.J. & Heath, M.R. (1997) Seasonal
 415 variation in the consumption of food by fish in the North Sea and implications for food web
 416 dynamics. *ICES Journal of Marine Science*, **54**, 243–266.

417 Hanchet, S. (1991) Diet of spiny dogfish, *Squalus acanthias* Linnaeus, on the east coast, South
 418 Island, New Zealand. *Journal of Fish Biology*, **39**, 313–323.

419 Hedger, R., McKenzie, E., Heath, M., Wright, P., Scott, B., Gallego, A., *et al.* (2004) Analysis of
 420 the spatial distributions of mature cod (*Gadus morhua*) and haddock (*Melanogrammus*
 421 *aeglefinus*) abundance in the North Sea (1980–1999) using generalised additive models.
 422 *Fisheries Research*, **70**, 17–25.

423 Holden, M.J. (1966) The Food of the Spurdog, *Squalus acanthias* (L.). *ICES Journal of Marine*
 424 *Science*, **30**, 255–266.

425 Hutchings, J.A. (2000) Collapse and Recovery of Marine Fishes. *Nature*, **406**, 882–885.

426 Hyslop E. J. (1980) Stomach contents analysis-a review of methods and their application.
 427 *Journal of Fish Biology*, **17**, 411–429.

428 International Council for the Exploration of the Seas (ICES). (1989) *ICES Cooperative Research*
 429 *Report No. 164*. Copenhagen, Denmark.

430 International Council for the Exploration of the Seas (ICES). (1997) *ICES Database report of the*
 431 *stomach sampling project 1991 No. 219*. Copenhagen, Denmark.

432 International Council for the Exploration of the Seas (ICES). (2010) *ICES Stomach Dataset*.
 433 Copenhagen, Denmark.

434 Jaworski, a & Ragnarsson, S. (2006) Feeding habits of demersal fish in Icelandic waters: a
 435 multivariate approach. *ICES Journal of Marine Science*, **63**, 1682–1694.

436 Jensen, A.C. (1965) *Fishery Leaflet 578: Haddock*. Washington, DC.

437 Jiang, W. & Jørgensen, T. (1996) The diet of haddock (*Melanogrammus aeglefinus* L .) in the
 438 Barents Sea during the period 1984 – 1991. *ICES Journal of Marine Science*, **53**, 11–21.

439 Jones, B.C. & Geen, G.H. (1977) Food and Feeding of Spiny Dogfish (*Squalus acanthias*) in
 440 British Columbia Waters. *Journal of Fisheries Research*, **34**, 2067–2078.

441 Kelly, C., Codling, E. & Rogan, E. (2006) The Irish Sea cod recovery plan: some lessons
 442 learned. *ICES Journal of Marine Science*, **63**, 600–610.

443 Lauzon-Guay, J.-S. & Scheibling, R.E. (2007) Behaviour of sea urchin *Strongylocentrotus*
 444 *droebachiensis* grazing fronts: food-mediated aggregation and density-dependent facilitation.
 445 *Marine Ecology Progress Series*, **329**, 191–204.

446 Link, J.S. (2004) Using fish stomachs as samplers of the benthos: integrating long-term and
 447 broad scales. *Marine Ecology Progress Series*, **269**, 265–275.

448 Link, J.S. (2007) Underappreciated Species in Ecology: “Ugly Fish” in the Northwest Atlantic
 449 Ocean. *Ecological Applications*, **17**, 2037–2060.

450 Link, J.S. (2011) Ecological Considerations in Fisheries Management: When Does It Matter?
 451 *Fisheries*, **27**, 10–17.

452 Link, J.S. & Auster, P.J. (2013) The Challenges of Evaluating Competition Among Marine
 453 Fishes: Who Cares, When Does It Matter, and What Can One Do About It? *Bulletin of Marine*

454 *Science*, **89**, 213–247.

455 Link, J.S., Bogstad, B., Sparholt, H. & Lilly, G.R. (2009) Trophic role of Atlantic cod in the
 456 ecosystem. *Fish and Fisheries*, **10**, 58–87.

457 Link, J.S. & Garrison, L.P. (2002a) Changes in piscivory associated with fishing induced
 458 changes to the finfish community on Georges Bank. *Fisheries Research*, **55**, 71–86.

459 Link, J.S. & Garrison, L.P. (2002b) Trophic ecology of Atlantic cod *Gadus morhua* on the
 460 northeast US continental shelf. *Marine Ecology Progress Series*, **227**, 109–123.

461 Link, J.S., Garrison, L.P. & Almeida, F.P. (2002) Ecological Interactions between
 462 Elasmobranchs and Groundfish Species on the Northeastern U. S. Continental Shelf .I.
 463 Evaluating Predation. *Journal of Fisheries Management*, **22**, 550–562.

464 Løkkeborg, S., Bjordal, A. & Fernö, A. (1989) Responses of Cod (*Gadus morhua*) and haddock
 465 (*Melanogrammus aeglefinus*) to baited hooks in the natural environment. *Canadian Journal of*
 466 *Fish and Aquatic Sciences*, **46**, 1478–1483.

467 MacDonald, J.S., Waiwood, K.G. & Green, R.H. (1982) Rates of digestion of different prey in
 468 Atlantic Cod (*Gadus morhua*), Ocean Pout (*Macrozoarces americanus*), Winter Flounder
 469 (*Pseudopleuronectes americanus*), and American Plaice (*Hippoglossoides platessoides*).
 470 *Canadian Journal of Fisheries and Aquatic Sciences*, **39**, 651–659.

471 Mahon, R. & Neilson, J.D. (1987) Diet changes in Scotian Shelf haddock during the pelagic and
 472 demersal phases of the first year of life. *Marine Ecology Progress Series*, **37**, 123–130.

473 Mattson, S. (1992) Food and feeding habits of fish species over a soft sublittoral bottom in the
 474 Northeast Atlantic. 3. Haddock (*Melanogrammus aeglefinus* (L.))(Gadidae). *Sarsia: North*
 475 *Atlantic Marine Science*, **77**, 33–45.

476 McNamara, J.M. & Houston, A.I. (1987) Partial preferences and foraging. *Animal Behaviour*,
 477 **35**, 1084–1099.

478 Moran, P.J., De'ath, G., Baker, V.J. & Bass, D.K. (1992) Pattern of Outbreaks of Crown-of-
 479 thorns Starfish (*Acanthaster planci* L.) along the Great Barrier Reef since 1966. *Australian*
 480 *Journal of Marine and Freshwater Research*, **43**, 555–568.

481 Murawski, S.A., Brown, R., Lai, H.-L., Rago, P.J. & Hendrickson, L. (2000) Large-Scale Closed
 482 Areas as a Fishery-Management Tool in Temperate Marine Systems: the Georges Bank
 483 Experience. *Bulletin of Marine Science*, **66**, 775–798.

484 Northeast Fisheries Science Center (NEFSC). (1988) *An evaluation of the bottom trawl survey*
 485 *program of the Northeast Fisheries Science Center. NOAA Technical Memorandum*. Woods
 486 Hole, MA. USA.

487 Northeast Fisheries Science Center (NEFSC). (1998) *Status of the fishery resources off the*
 488 *Northeastern United States from 1998. NOAA Technical Memorandum*. Woods Hole, MA. USA.

489 Packer, D.B., Watling, L. & Langton, W. (1994) The population structure of the brittle star
 490 *Ophiura sarsi* Lotken in the Gulf of Maine and its trophic relationship to American plaice
 491 (*Hippoglossoides platessoides* Fabricius). *Journal of Experimental Marine Biology and Ecology*,
 492 **179**, 207–222.

493 Paine, R.T. (1966) Food web complexity and species diversity. *American Naturalist*, **100**, 65–75.

494 Paine, R.T. (1969) The *Pisaster-Tegula* interaction: prey patches, predator food preference and
 495 intertidal community structure. *Ecology*, **50**, 950–961.

496 Pauly, D., Christensen, V., Dalsgaard, J., Froese, R., Jr, F.T., Series, N., *et al.* (1998) Fishing
 497 down marine food webs. *Science*, **279**, 860–863.

498 Pecuchet, L., Nielsen, J.R. & Christensen, A. (2014) Impacts of the local environment on
 499 recruitment: a comparative study of North Sea and Baltic Sea fish stocks. *ICES Journal of*
 500 *Marine Science*.
 501 Perry, R.I. & Smith, S.J. (1994) Identifying habitat associations of marine fishes using survey
 502 data: an application to the northwest Atlantic. *Canadian Journal of Fisheries and Aquatic*
 503 *Sciences*, **51**, 589–602.
 504 Pinnegar, J.K. (2014) DAPSTOM - An integrated database and portal for fish stomach records.
 505 Version 4.7. [WWW Document]. *Centre for the Environment, Fisheries & Aquaculture Science*.
 506 URL <http://www.cefas.defra.gov.uk/our-science/fisheries-information/fish-stomach-records.aspx>
 507 [accessed on 2014].
 508 Pinnegar, J.K., Polunin, N.V.C., Francour, P., Badalamenti, F., Chemello, R., Harmelin-Vivien,
 509 M.-L., *et al.* (2000) Trophic cascades in benthic marine ecosystems: lessons for fisheries and
 510 protected-area management. *Environmental Conservation*, **27**, 179–200.
 511 Platt, T., Fuentes-Yaco, C. & Frank, K.T. (2003) Marine ecology: spring algal bloom and larval
 512 fish survival. *Nature*, **423**, 398–399.
 513 Pope, J.G. & Macer, C.T. (1996) An evaluation of the stock structure of North Sea cod, haddock,
 514 and whiting since 1920, together with a consideration of the impacts of fisheries and predation
 515 effects on their biomass and recruitment. *ICES Journal of Marine Science*, **53**, 1157–1169.
 516 Rowlands, W.L., Dickey-Collas, M., Geffen, A.J. & Nash, R.D.M. (2008) Diet overlap and prey
 517 selection through metamorphosis in Irish Sea cod (*Gadus morhua*), haddock (*Melanogrammus*
 518 *aeglefinus*), and whiting (*Merlangius merlangus*). *Canadian Journal of Fisheries and Aquatic*
 519 *Sciences*, **65**, 1297–1306.

520 Schückel, S., Ehrich, S., Kröncke, I. & Reiss, H. (2010) Linking prey composition of haddock
 521 *Melanogrammus aeglefinus* to benthic prey availability in three different areas of the northern
 522 North Sea. *Journal of Fish Biology*, **77**, 98–118.

523 Sell, A.F. & Kröncke, I. (2013) Correlations between benthic habitats and demersal fish
 524 assemblages — A case study on the Dogger Bank (North Sea). *Journal of Sea Research*, **80**, 12–
 525 24.

526 Shackell, N.L., Frank, K.T., Fisher, J.A.D., Petrie, B. & Leggett, W.C. (2010) Decline in top
 527 predator body size and changing climate alter trophic structure in an oceanic ecosystem.
 528 *Proceedings of the Royal Society B*, **277**, 1353–60.

529 Skold, M., Loo, L. & Rosenberg, R. (1994) Production, dynamics and demography of an
 530 *Amphiura filiformis* population. *Marine Ecology Progress Series*, **103**, 81–90.

531 Smith, B., Collie, J. & Lengyel, N. (2013) Effects of chronic bottom fishing on the benthic
 532 epifauna and diets of demersal fishes on northern Georges Bank. *Marine Ecology Progress*
 533 *Series*, **472**, 199–217.

534 Smith, B.E., Ligenza, T.J., Almeida, F.P. & Link, J.S. (2007) The trophic ecology of Atlantic
 535 cod: insights from tri-monthly, localized scales of sampling. *Journal of Fish Biology*, **71**, 749–
 536 762.

537 Smith, B.E. & Link, J.S. (2010) *The Trophic Dynamics of 50 Finfish and 2 Squid Species on the*
 538 *Northeast US Continental Shelf*. Woods Hole, MA. USA.

539 Sonina, M.A. (1969) Biology of arcto-Norwegian haddock during 1927-1965. *Firsheries*
 540 *Research Board of Canada*, **1924**, 1–154.

541 Steimle, F.W. (1990) *Benthic Macrofauna and Habitat Monitoring on the Continental Shelf of*

542 *the Northeastern United States*. Washington, DC. USA.

543 Steimle, F.W. & Terranova, R.J. (1980) Energy Equivalents of Marine Organisms from the
 544 Continental Shelf of the Temperate Northwest Atlantic. *Journal of the Northwest Atlantic*
 545 *Fishery Science*, **6**, 117–124.

546 Tanasichuk, R.W., Ware, D.M., Shaw, W. & Mcfarlane, G.A. (1991) Variations in diet, daily
 547 ration, and feeding periodicity of Pacific Hake (*Merluccius productus*) and Spiny Dogfish
 548 (*Squalus acanthias*) off the lower west coast of Vancouver Island. *Canadian Journal of Fish and*
 549 *Aquatic Sciences*, **48**, 2118–2128.

550 Temming, A., Gotz, S., Mergardt, N. & Ehrich, S. (2004) Predation of whiting and haddock on
 551 sandeel: aggregative response, competition and diel periodicity. *Journal of Fish Biology*, **64**,
 552 1351–1372.

553 Tibbetts, S.M., Lall, S.P. & Milley, J.E. (2004) Apparent digestibility of common feed
 554 ingredients by juvenile haddock, *Melanogrammus aeglefinus* L. *Aquaculture Research*, **35**, 643–
 555 651.

556 Tibbetts, S.M., Milley, J.E. & Lall, S.P. (2006) Apparent protein and energy digestibility of
 557 common and alternative feed ingredients by Atlantic cod , *Gadus morhua* (Linnaeus , 1758).
 558 *Aquaculture*, **261**, 1314–1327.

559 Tollit, D.J., Black, a. D., Thompson, P.M., Mackay, A., Corpe, H.M., Wilson, B., *et al.* (1998)
 560 Variations in harbour seal *Phoca vitulina* diet and dive-depths in relation to foraging habitat.
 561 *Journal of Zoology*, **244**, 209–222.

562 Tseeb, R.Y. (1964) *Feeding and food adaptations of haddock*. Nauka, Moscow, St. Petersburg.

563 Vinyard, G.L. (1980) Differential prey vulnerability and predator selectivity : effects of evasive

564 prey on Bluegill (*Lepomis macrochirus*) and Pumpkinseed (*L. gibbosus*) predation. *Canadian*
565 *Journal of Fish and Aquatic Sciences*, **37**, 2294–2299.

566 Vogler, C., Benzie, J., Lessios, H., Barber, P. & Wörheide, G. (2008) A threat to coral reefs
567 multiplied? Four species of crown-of-thorns starfish. *Biology letters*, **4**, 696–9.

568 Watanabe, J.M. & Harrold, C. (1991) Destructive grazing by sea urchins *Strongylocentrotus* spp.
569 in a central California kelp forest: potential roles of recruitment, depth and predation. *Marine*
570 *Ecology Progress Series*, **71**, 125–141.

571 Wood, H.L., Widdicombe, S. & Spicer, J.I. (2009) The influence of hypercapnia and the infaunal
572 brittlestar *Amphiura filiformis* on sediment nutrient flux – will ocean acidification affect nutrient
573 exchange? *Biogeosciences*, **6**, 2015–2024.

574 Zamarro, J. (1992) Feeding behaviour of the American Plaice (*Hippoglossoides platessoides* on
575 the Southern Grand Bank of Newfoundland. *Netherlands Journal of Sea Research*, **29**, 229–238.

576

577 Table I. Data collection methods and sources. Frequency of occurrence was measured as the number of each diet item relative to the
578 total number of stomachs in percent. Diet composition was measured as the weight of the diet items relative to the total weight of *M.*
579 *aeglefinus* stomachs in percent.

Ecosystem	Time of sampling	Type of data collected	<i>M. aeglefinus</i> stomachs sampled (#)	Collection method	Source
Gulf of Maine	1973-1982, 1984-2011	Diet composition	3629	Data compiled from Northeast Fisheries Science Center surveys through random stratified bottom trawl surveys.	(Northeast Fisheries Science Center (NEFSC), 1988; Smith & Link, 2010)
Georges Bank	1973-1982, 1984-1992, 1994-2011	Diet composition	5825	Data compiled from Northeast Fisheries Science Center surveys through random stratified bottom trawl surveys.	(Northeast Fisheries Science Center (NEFSC), 1988, 1998; Smith & Link, 2010)
Scotian Shelf	1995-2009	Diet composition	2690	Data compiled from Department of Fisheries and Oceans trawl surveys.	(Cook & Bundy, 2010, 2012; Bundy <i>et al.</i> , 2011)
North Sea	1981, 1991	Diet composition	4452	Data collected during the quarterly International Bottom Trawl Surveys of the North Sea	(International Council for the Exploration of the Seas (ICES), 1989, 1997, 2010)
Rockall Bank	2004-2007, 2009-2010	Diet composition	3746	Data compiled from the Polar Research Institute of Marine Fisheries and Oceanography.	(Antipova <i>et al.</i> , 1980)
Barents Sea	1984-1991, 1993-2011	Diet composition	11064	Data compiled by the Institute of Marine Research and the Polar Research Institute of Marine Fisheries and Oceanography.	(Antipova <i>et al.</i> , 1980; Dolgov <i>et al.</i> , 2007)
Irish Sea	1894, 1981-1983, 2008, 2010	Frequency of occurrence	653	Data compiled from the Centre for Environment, Fisheries and Aquaculture Science.	(Pinnegar, 2014)
Celtic Sea	1977, 1978, 1981-1988, 1991-1993	Frequency of occurrence	971	Data compiled from the Centre for Environment, Fisheries and Aquaculture Science.	(Pinnegar, 2014)
Spitzbergen	1930, 1936, 1937, 1949, 1950, 1954, 1964	Frequency of occurrence	735	Data compiled from the Centre for Environment, Fisheries and Aquaculture Science.	(Pinnegar, 2014)

580 Table II. Results of the Chi-square analysis examining the relationship between echinoderms and fish in *M. aeglefinus* diet within each
 581 ecosystem. Bold numbers indicate significant differences.

Ecosystem	X^2	p -value	Dominant prey
Gulf of Maine	3.79	0.05	None
Georges Bank	0.12	0.73	None
Scotian Shelf	2.80	0.09	None
North Sea	6.92	< 0.01	Fish
Rockall Bank	22.11	< 0.01	Fish
Barents Sea	0.02	0.89	None
Irish Sea	3.77	0.05	None
Celtic Sea	22.54	< 0.01	Echinoderms
Spitzbergen	43.67	< 0.01	Echinoderms

582

583

584

585

586

587

588

589

590 Table III. Results of the Chi-square analysis (p-values) examining the relationship between echinoderms and fish in *M. aeglefinus* diet
 591 at each five year block within ecosystems. Bold numbers indicate significant differences.

Year	Georges Bank	Gulf of Maine	Scotian Shelf	Barents Sea
1975	< 0.01	< 0.01		
1980	< 0.01	< 0.01		
1985	< 0.01	< 0.01		0.03
1990	< 0.01	< 0.01		0.90
1995	0.08	< 0.01		< 0.01
2000	< 0.01	< 0.01	< 0.01	0.85
2005	< 0.01	0.82	< 0.01	0.09
2010	0.29	0.11	< 0.01	0.10
2015	0.56	0.71		0.59


FIGURE CAPTIONS

Figure 1. Map of different ecosystems sampled.

Figure 2. Percent contribution of echinoderms and fish to *M. aeglefinus* from different ecosystems. Diet from Celtic Sea, Irish Sea and Spitzbergen were measured in frequency of occurrence (number of stomachs containing the diet item relative to the total number of stomachs), diet from the Barents Sea, North Sea, Rockall Bank, Scotian Shelf, Georges Bank and Gulf of Maine was measured in diet composition (weight of the diet items relative to the total weight of *M. aeglefinus* stomachs in percent).

Figure 3. Percent contribution of echinoderms and fish to the diet of different size classes (cm) of *M. aeglefinus* in different ecosystems where size class data was available.

Figure 4. Percent contribution of echinoderm (grey dotted line) and fish (black solid line) over time (year) in different ecosystems where time series data was available.



616

617 Figure 1. Map of different ecosystems sampled.

618




Figure 2. Diet of *M. aeglefinus* in different ecosystems.





Figure 3. Diet of different size class of *M. aeglefinus*.



626

627 Figure 4. Diet of *M. aeglefinus* over time.